

## PHENOLIC COMPOSITION, ANTIMICROBIAL ACTIVITY OF *Rosmarinus officinalis*

Reçu le 27/02/2014 – Accepté le 16/09/2014

S. ATHAMENA<sup>1</sup>, S. LAROUI<sup>1</sup>, M. ATHAMENA<sup>2</sup>

<sup>1</sup> Biotechnology Laboratory of bioactive molecules and cellular pathophysiology, University of Batna, Algeria

<sup>2</sup> Service of internal medicine CHU, Faculty of Medicine, University of Batna, Algeria

Email : athamenasouad@yahoo.fr

### Résumé

L'objectif du travail est l'évaluation de l'activité antimicrobienne des extraits du romarin (EBr, EAcOEt et En-BuOH) sur huit souches bactériennes (*Escherichia coli* ATCC, *Pseudomonas aeruginosa* ATCC, *Staphylococcus aureus* ATCC deux souches d'*Enterobacter* sp, *Serratia* sp, *Klebsiella pneumoniae* et *Streptocoque* sp) et trois souches fongiques (*Candida albicans*, *Candida Kefyr* et *Aspergillus niger*) par la méthode de diffusion en milieu gélosé. Les dosages quantitatifs des polyphénols totaux par la méthode de Folin-Ciocalteu ont révélé la richesse du romarin en polyphénols ( $195,45 \pm 4,16$  mg EAG/g d'EBr). L'analyse qualitative par HPLC a révélé la présence de la rutine et la catéchine dans les extraits du romarin. Les extraits du romarin ont présenté une activité antibactérienne qui s'étend sur la totalité des souches de la collection et l'extrait d'acétate d'éthyle (EAcOEt) s'est révélé le plus actif et témoigne d'une activité antibactérienne intéressante surtout contre la bactérie multi-résistante *Enterobacter* sp (BLSE+CHN). Seule l'inhibition de la souche *Klebsiela pneumoniae* a présenté une corrélation très significative entre le taux des polyphénols de l'EBr du romarin et l'activité antibactérienne ( $R^2 = 0,992$ ). Les extraits du romarin se sont révélés inactifs vis-à-vis les souches fongiques.

**Mots clés :** *Rosmarinus officinalis*, Polyphénols, Activité antimicrobienne.

### Abstract

This work aims at evaluating the antimicrobial activity of rosemary extracts (EBr, EAcOEt and En-BuOH) on 8 strains of bacteria (*Escherichia coli* ATCC, *Pseudomonas aeruginosa* ATCC, *Staphylococcus aureus* ATCC 2 strains of *Enterobacter* sp, *Serratia* sp, *Klebsiella pneumoniae* and *Streptococcus* sp) and 3 fungal strains (*Candida albicans*, *Candida Kefyr* and *Aspergillus niger*), by agar diffusion method. The quantification of total polyphenols using the Folin-Ciocalteu method and of the flavonoids revealed the richness of the rosemary in polyphenols ( $195,45 \pm 4,16$  mg EAG/g of EBr). The analysis by HPLC revealed the presence of rutin and the catechin in the extracts rosemary. The results revealed that the extracts of rosemary are showed antibacterial activity against the whole tested Bacterial strains and The rosemary EAcOEt has been the most active extract and it has revealed an interesting antibacterial activity against the multi-resistant strain *Enterobacter* sp (ESBL+HLC). Only inhibition of *Klebsiela pneumonia* strain have shown very significatif correlation between polyphenolic content of EBr of rosemary and antibacterial activity ( $R^2 = 0,992$ ). The results of the antifungal activity showed the inefficiency of all the extracts against the fungal strains.

**Key words:** *Rosmarinus officinalis*, phenolic compounds, Antimicrobial activity.

### ملخص

إن الهدف من هذه الدراسة هو اختبار التأثير النشاط المضاد للميكروبات ضد ثمان سلالات بكتيرية : *Escherichia coli* : ATCC, *Pseudomonas aeruginosa* ATCC, *Staphylococcus aureus* ATCC 2 souches d'*Enterobacter* sp, *Candida albicans*, *Candida Kefyr* et *Aspergillus niger* عن طريق الانتشار على وسط صلب: لمستخلصات (المستخلص الخام: مستخلص خلات الإثيل ومستخلص البوتانول العادي) لأوراق نبتة الإكليل. فمنا أولا بإجراء تقدير كمي للفينولات و كذلك الفلافونيدات على أساس أنها أهم قسم من العائلة الفينولية و دراسة تحليية للفلافونيدات بواسطة كروماتوغرافيا السائل العالي الأداء (HPLC). التقدير الكمي للفينولات بواسطة طريقة Folin-Ciocalteu يبيّن غنّى الإكليل بالفينولات ( $195,45 \pm 4,16$  mg EAG/g d'EBr) (الدراسة التحليلية بواسطة HPLC يبيّن وجود rutine و catéchine في كل في مست لصات الإكليل. مستخلص خلات الإثيل لنبتة الإكليل كان الأكثر فعالية على مجموع البكتيريات المختبرة و أظهر نشاطاً مهماً مضاداً للبكتيريا ضد السلالة المقاومة *Enterobacter* sp (BLSE+CHN). تثبيط السلالة *Klebsiela pneumonia* هو الوحيد الذي اظهر ارتباط دو دلالة إحصائية بين المحتوى الفينولي مستخلص الإكليل الخام و النشاط المضاد للبكتيريا ( $R^2 = 0,992$ ). نتائج النشاط المضاد للبكتيريات يبيّن عدم فعالية كل المستخلصات ضد السلالات الفطرية.

**الكلمات المفتاحية :** الإكليل، الفينولات، النشاط المضاد للميكروبات.



In recent years, there has been a great interest for the discovery of new antimicrobial agents, due to an alarming increase in the rate of infections with micro-organisms resistant to antibiotics. One of the common approaches to the search for biologically active substances is the systematic screening of micro-organisms or plants, which are a source of many useful therapeutic agents.

The antimicrobial activity of oils and extracts of plants have formed, in particular, the basis for many applications, including pharmaceutical, medicine, natural therapy and food conservation [1].

The plant matter contains a large number of molecules that have multiple interests used in the industry, food, cosmetology and dermopharmacy. Among these compounds, we can find coumarins, alkaloids, phenolic acids, tannins, lignans, terpenes and flavonoids [2].

The rosemary (*Rosmarinus officinalis L.*) is the object of recent research in the fields of pharmaceuticals, cosmetics and food industry. It is an aromatic grass which is presented in the form of shrub, under sapling or herbaceous that belongs to the family of *Labiees* [3], measuring approximately 0.8 to 2m in height [4].

The leaves are closely linear lanceolate, brittle and tough. The flowers of a pale blue, stained inwardly with purple are arranged in short dense clusters flourish throughout most of the year.

The rosemary is very appreciated for its aromatic properties, anti-oxidant, antimicrobial, antispasmodic, emmenagogues and anti-tumor, widely used in the pharmaceutical products and in traditional medicine [3].

The aim of this work is to evaluate the antimicrobial activity by the method of diffusion in an agar medium of crude extract and their fractions (EAcOEt, En-BuOH) of the medicinal plant, the rosemary. This assessment is linked to the phenolic content of these extracts.

## MATERIAL AND METHODS

### Plant Material and preparation of extracts

The extraction of flavonoids is carried out according to the diagram presented by Lebreton (1967) as amended by Boutard (1972), Gonnet (1973) and Jay (1975).

The rosemary dry leaves are left to macerate overnight at ambient temperature, in a water-alcohol mixture of methanol-water (7:3 V/V). After filtration, the solvent is removed from the filtrate by rotary evaporation. The crude extract obtained is subjected to a liquid-liquid extraction successively by two solvents (ethyl acetate and the n- butanol). The series of extraction enabled us to obtain four fractions ; the crude extract hydro-methanolic (EBr), the fraction of ethyl acetate (EAcOEt), the fraction of the n-butanol (En-B OH) and the aqueous fraction (AqE) residual.

### Total phenolics compound Content

The total phenolic compound content was carried out with the colorimetric reagent Folin-Ciocalteu according to the method quoted by Wong [5]. 200 $\mu$ l of each extract were added to 1ml of 1:10 diluted Folin-Ciocalteu reagent. The solutions were mixed and incubated for 4 minutes.

After incubation, 800  $\mu$ l of a solution of sodium carbonate Na<sub>2</sub>CO<sub>3</sub> (75 g /l) have been added. The final mixture is incubated for 2 hours in the dark at ambient temperature. The absorbance was measured by a spectrophotometer at 765 nm. The content in polyphenols is expressed in milligram of gallic acid equivalents per gram of extract ( $\mu$ g EAG/mg).

### Total flavonoids Content

The method of aluminum trichloride (AlCl<sub>3</sub>) [6] is used to quantify the flavonoids in our extracts. 1 ml of each extract was added to an equal volume of a solution of AlCl<sub>3</sub> (2%). The mixture was vigorously agitated and the absorbance at 430 nm was read after 10 minutes of incubation. The results are expressed in milligrams of quercetin equivalent per gram of extract ( $\mu$ g EQ/mg).

## Qualitative analysis by HPLC

The analysis is carried out by an HPLC (VP SHIMADZU LIQUID CHROMATOGRAPH). 20 µl of each extract were injected on a column of type reverse phase C18, of equal size to 125 x 4.6 mm. The mobile phase consists of three eluents: distilled water, methanol, acetic acid (50: 47: 2.5) (V /V /V). The elution gradient applied is of an isocratic type spread over 10 min. The flow rate is 1 ml/min [7]. The detection was performed by a UV-Vis detector at a wavelength equal 254 nm.

The flavonoids in each extract analysis have been identified by the comparison of the retention times obtained by those witnesses.

## Antimicrobial Activity

The antibacterial activity of the extracts was determined by the agar diffusion method standardized by (NCLLS) [8].

Eight (n = 8) Bacterial strains have been tested :

1. *Escherichia coli* ATCC 25922,
2. *Pseudomonas aeruginosa* ATCC 27853,
3. *Staphylococcus aureus* ATCC 25923,
4. *Enterobacter* sp (HLC+ ESBL),
5. *Enterobacter* sp,
6. *Serratia* sp,
7. *Klebsiella pneumoniae* sp (ESBL),
8. *Streptococcus* sp.

Three (n = 3) Fungal strains: Two yeasts: *Candida albicans*, *Candida Kefyr* and a fungus: *Aspergillus niger*.

Of microbial communities well isolated were transferred into tubes of sterile distilled water in order to have a microbial suspension having a cell density adjacent to that of Mc Farland 0.5 (10<sup>6</sup> CFU/ml). It was subsequently spread the entire surface of the Agar Agar (Mueller Hinton for the bacteria non-demanding; Agar Mueller Hinton, which contains 5% blood of the horse for the bacteria demanding; Sabouraud for yeasts) by the microbial suspension.

For the fungus the Sabouraud agar is inoculated by the suspension.

The disks sterile impregnated of increasing concentrations of extracts resumed with the Dimethyl Sulfoxide (DMSO) to reason of 10 µl per disc [9] have been deposited out by sterile methods on the agar surface. The boxes have been incubated 24 h at 37 °C in normal atmosphere for the bacteria non-demanding and in an atmosphere containing 5% CO<sub>2</sub> for the bacteria demanding. The yeasts were incubated 48 h at 37 °C, while the fungus has been incubated 10 days at 27 °C in normal atmosphere. The antibacterial activity was expressed by measuring the diameter of the inhibition zone.

## Statistical study

The statistical study has been carried out by the statistical software Graph Pad Prism. All experiments were performed in triplicate; the results are expressed in average ± SD.

## RESULTS AND DISCUSSION

### Content in polyphenols

Generally, all plants of the family *Lamiaceae* are known for their phenolic compounds ([10], [11]). This is in accordance with our results presented in the Table 1.

**Table 1:** Polyphenols content of rosemary extracts.

Extract	Polyphenols contents <sup>(a)</sup>
EBr	195.45 ± 4.16
EAcOEt	541.82 ± 3.15
En-BuOH	539.39 ± 5.25

(a) mg of equivalent of gallic acid per gram of extract.

The values represent the average of 3 measurements ± SD

The content of CE rosemary is so close to that of Erkan *et al.* , [12] : 162 mg GAE/g and Ho *et al.* , [13] : 127 ± 3 mg GAE /g, but far enough to that of Tsai *et al.* , [14] : 58.1 ± 0.9 mg GAE/g and Tawaha *et al.* , [15] : 39.1 ± 3.6 mg GAE/g.

The shifted results result likely of:

The low specificity of the Folin-Ciocalteu reagent is the primary disadvantage of the colorimetric assay. The reagent is extremely sensitive to the reduction of all the groups of hydroxyl, not only those of phenolic compounds, but also of certain sugars and proteins etc. [16].

The distribution of secondary metabolites can change during the development of the plant. This may be linked to the harsh climate conditions (high temperature, solar exposure, drought, salinity), which stimulate the biosynthesis of secondary metabolites such as polyphenols [17].

### Content in flavonoids

The main reason for the choice of this class of polyphenols, lies in the fact that the flavonoids represent the most important polyphenols class, with more than 5000 compounds already described [16].

The results of the flavonoids assay are represented in table 2.

**Table 2:** Flavonoids Content rosemary extracts

Extract	Flavonoids Content <sup>(b)</sup>
EBr	2.06 ± 1.14
EAcOEt	21.39 ± 0.72
En-BuOH	19.58 ± 2.75

(b) mg of equivalent of quercetin per gram of extract. The values represent the average of 3 measurements ± SD

According to the results of Ho and his collaborators [13] the extract of rosemary is rich in polyphenols ( $127 \pm 3$  mg EAG/g) and poor in flavonoids ( $20.1 \pm 1.30$  mg EC/g). We can say that our results confirm those of Ho.

In addition, Tsai *et al.*, [14] have also found that the methanolic extract of the rosemary contains  $60.7 \pm 1.1$  mg EC/g. The levels reported by Ho and Tsai are very high compared with our results; this difference can probably explained by the difference of the standard used for the assay of the flavonoids.


Maisuthisakul *et al.*, [18] have found that the total content of the ethanolic flavonoids extracts of 28 plants, is linked to the content of the total phenolic compounds. Similarly, we have found that the content of the flavonoids extracts of the

rosemary was correlated significantly with the content of polyphenols ( $R^2 = 0.969$ ).


### Qualitative analysis by HPLC

The chromatograms of HPLC of the different extracts are represented below.


**Figure 1:** Chromatogram of HPLC of EBr of rosemary registered to 254 nm



**Figure 2:** Chromatogram of HPLC of EAcOEt rosemary recorded at 254 nm



**Figure 3:** Chromatogram of HPLC of En-BuOH rosemary recorded at 254 nm



The comparison of the retention times (Table 3) of the standards with those recorded in the different chromatograms (Table 4), allows a possible identification of some flavonoids in our extracts [19].

**Table 3:** Retention time of the flavonoids standards.

Retention Time (min)	The Standard flavonoid
1.8	Quercetin
3.4	Rutin
2.0	Catechin

**Table 4:** Retention time of the flavonoids present in the rosemary extracts.

Retention Time (min)				The possible flavonoid
Ebr	EAcOEt	En-	BuOH	
1.7	0.5	0.8	-	
2.0	0.8	1.7	Presence of the catechin in the EBr	
2.6	1.2	2.0	Presence of the catechin En-BuOH	
3.4	1.7	3.4	Presence of the rutin in the Ebr and En-BuOH	
3.8	2.0	3.7	Presence of the catechin in the EAcOEt	
4.7	2.6	4.7	-	
5.2	3.4	5.3	Presence of the rutin in the EAcOEt	
6.3	3.7	7.5	-	
7.6	4.6	9.4	-	
8.9	5.5			
9.6	6.1			
	7.5			
	8.0			
	8.8			

The results show the presence of the catechin, the rutin and the absence of quercetin in the totality of the extracts.

Similarly Justesen *et al.*, [20] and Wojdylo *et al.*, [21] who have used as a mobile phase a gradient system have revealed the absence of quercetin in the methanolic rosemary extracts.

## Antimicrobial Activity

The results presented in the tables below show that:

- The rosemary extracts display important activities, which extend on the totality of the collection strains, including the EAcOEt which is the most active.
- The strains *Escherichia coli* ATCC and *Klebsiella pneumoniae* possess a very high resistance potential against the antibacterial action of 3 rosemary extracts.
- A few zones of inhibition with moderate the EAcOEt were recorded with *Pseudomonas aeruginosa* ATCC ( $11.33 \pm 1.15$ mm), the two strains of *Enterobacter* sp ( $13.00 \pm 0.00$  ; AT  $15.67 \pm 0.58$  ) and *Serratia* sp ( $12.33 \pm 1.53$ mm).
- The *Enterobacter* sp (ESBL+HLC) is a bacterium highly resistant to antibiotics, but has proved to be very sensitive to the EAcOEt rosemary than the bacterium *Enterobacter* sp sensitive to Cefotaxim.
- A higher activity with the 3 rosemary extracts, has been noticed in *Staphylococcus aureus* ATCC, which is sensitive to the low concentrations of extracts.
- *Sterptocoque* sp, a bacterium Gram (+), has proved resistant to extracts tested.
- The inhibitor effects increase considerably with the concentration of extracts. The majority of extracts can retain a detectable activity, after weak dilutions. The Rosemary EAcOEt has remained active also after the 1/16 dilution for strains *Pseudomonas aeruginosa* ATCC, *Staphylococcus aureus* ATCC and two strains of *Enterobacter* sp.

**Table 5:** Diameter of the inhibition zone of the rosemary EBr.

Bacterial strains	Diameter of the inhibition zone * (mm)			
	The dilutions of the EBr rosemary			
	1/2	1/4	1/8	1/1
				6
<i>Escherichia coli</i> ATCC	-	-	-	-
<i>Pseudomonas aeruginosa</i> ATCC	10.00 ± 1.73	8.67 ± 1.15	7.3 3	-
<i>Staphylococcus aureus</i> ATCC	25.33±1.15	23.67± 0.58	17.67 ± 0.5	17.33 ± 0.5
<i>Enterobacter</i> sp	9.00 ± 1.00	8.00 ± 0.00	-	-
<i>Enterobacter</i> sp BLSE+HLC	12.33 ± 1.15	9.66 ± 0.58	9.3 3	-
<i>Klebsiella pneumoniae</i> BLSE	7.67 ± 0.58	-	-	-
<i>Serratia</i> sp	7.00 ± 0.00	-	-	-
<i>Sterptocoque</i> sp	7.67 ± 0.58	-	-	-

(\*) Diameter of the inhibition zone produced around the disks by the addition of 10 µl of extract.  
(Diameter of the disc is included)

**Table 6:** Diameter of the inhibition zone of the rosemary EAcOEt.

Bacterial strains	Diameter of the inhibition zone * (mm)			
	The dilutions of the EAcOEt rosemary			
	1/2	1/4	1/8	1/16
<i>Escherichia coli</i> ATCC	8.00 ± 0.00	-	-	-
<i>Pseudomonas aeruginosa</i> ATCC	11.33 ± 1.15	11.33 ± 1.15	9.33 8	8.67 ± 0.5
<i>Staphylococcus aureus</i> ATCC	28.33 ± 0.58	27.67 ± 0.5	24.67 ± 0.5	23.33 ± 0.58
<i>Enterobacter</i> sp	13.00 ± .00	12.33 ± 0.58	9.33 1	9.33 ± 0.5
<i>Enterobacter</i> sp ESBL+HLC	15.67 ± 0.58	16 ± 2.0	13.67 0.58	11.00 ± 2.64
<i>Klebsiella pneumoniae</i> ESBL	-	-	-	-
<i>Serratia</i> sp	12.33 ± 1.53	12.33 ± 0.58	9.67 3	7.33 ± 1.1
<i>Sterptocoque</i> sp	8.67 ± 1.15	8.00 ± 0	-	-

(\*) Diameter of the inhibition zone produced around the disks by the addition of 10 µl of extract. (Diameter of the disc is included)

The results reveal variable answers in function of the strains, of the concentration, type of the tested extract and that the sensitivity or resistance to antibiotics has no relation with that of the extracts.

Several works have highlighted the great sensitivity of the bacteria Gram (+) compared to the Gram (-) ([17]; [22]; [23]; [24]; [25]), this can be attributed to the difference in the outer layers of bacteria Gram (-) and Gram (+).

The bacteria Gram (-), independently of the cells membrane, possess an additional layer: the outer membrane, which is composed of phospholipids, proteins and lipopolysaccharides. This membrane is impermeable to most molecules. Nevertheless, the presence of porins in

this layer allows the free diffusion of molecules with a molecular mass below 600 Da. However, the inhibition of the Gram (-) bacteria growth has been reported, particularly in combination with the factors that can disturb the integrity of the cell and/or permeability of the membrane, such as low values of pH and increased concentrations in NaCl [ 26].

**Table 7:** Diameter of the inhibition zone of the En-BuOH of rosemary.

Bacterial strains	Diameter of the inhibition zone * (mm)			
	The dilutions of the En-BuOH rosemary			
	1/2	1/4	1/8	1/16
<i>Escherichia coli</i> ATCC	-	-	-	-
<i>Pseudomonas aeruginosa</i> ATC C	-	-	-	-
<i>Staphylococcus aureus</i> ATCC	24.00 ± 0.0	24.0 0	19.6 ± 7	18.3 ± 3
<i>Enterobacter</i> sp	-	-	-	-
<i>Enterobacter</i> sp also	8.33 ± 0.58	-	-	-
<i>Klebsiella pneumoniae</i> ESBL	-	-	-	-
<i>Serratia</i> sp	-	-	-	-
<i>Sterptocoque</i> sp	9.67 ± 0.58	-	-	-

The hypersensitivity of the strain *Staphylococcus aureus* ATCC can be explained by the probability of the sensitivity of bacteria Gram (+) to external environmental changes, such as temperature, pH and the natural extracts due to the absence of the outer membrane [27]. Some studies show no selective antimicrobial activity towards the bacteria Gram (+) or Gram (-) [28]. The resistance of the strain *Sterptocoque* sp can be attributed to the ability of the antibacterial agent to diffuse uniformly in the agar [22].

The inhibition zone increases significantly with the concentration of the extracts, a fact also

noticed by Dordevic and his collaborators, [29]. The disk load affects the antimicrobial activity, Rasooli and his collaborators, [30] have noted that the inhibition of the growth of *Aspergillus parasiticus* is strong when the disk is more responsible in essential oils of *Rosmarinus officinalis* and *Trachyspermum Copticum*.

The method used for the evaluation of the antibacterial activity also affects the results Natarajan *et al.*, (31) and Fazeli *et al.*, (32) have found that the method of dissemination from wells on agar is more suitable for studying the activities of aqueous extracts and organic of the *Euphorbia fusicornis* and Hydro-ethanolics of *Rhus coriaria* and *Zataria multiflora*, than the method of agar diffusion.

Polyphenols, such as tannins and flavonoids like epigallocatechol, the catechin, the myricetin, quercetin, [24] and luteolin [33] are important antibacterial substance. The HPLC has revealed the presence of the catechin in all extracts of rosemary, which may explain the antibacterial activity of the extracts of this plant.

We have found that there is not a correlation between the content of rosemary phenolic extracts and antibacterial activity. The values of the correlation coefficient  $R^2$  calculated were between  $R^2 = 0,442$  and  $R^2 = 0.001$  ( $P < 0.05$ ) and decrease in the following order:

1. *Sterptococcus* sp ;
2. *Escherichia coli* ATCC ;
3. *Pseudomonas aeruginosa* ATCC;
4. *Staphylococcus aureus* ATCC,
5. *Enterobacter* sp
6. and *Enterobacter* sp ESBL+HLC.

Similar results ( $R^2 = 0.00$ ) were obtained by Turkmen *et al.*, [23] during the evaluation of the antibacterial activity of the extracts of tea.

Only the inhibition of the strain *Klebsiella pneumoniae* which presented a correlation very significantly, between the rate of polyphenols of the EBr of the rosemary and the antibacterial activity with a coefficient  $R^2 = 0,992$ . This result

is consistent with the first systematic study prepared by Shan *et al.*, [24].

In fact this study shows a report highly positive between antibacterial activity and the rate of the polyphenols of a large number of extracts (46) of spices and herbs, the values of the coefficients of correlation  $R^2$  were between 0.93 and 0.73.

The antimicrobial activity depends not only on the presence of phenolic compounds, but also on the presence of various secondary metabolites [34], on the location and on the number of hydroxyl groups [17].

Although, ethanol and methanol were the best solvents than other ones by extracting the phenolic compounds, because of their polarity and of their good solubility for these compounds, the results have proven that ethanol was the best solvent to extract the phenolic compounds, followed by methanol and finally by water [35] which could explain the difference mentioned below.

We found a difference by comparing the inhibition zone of *Escherichia coli* ATCC (0.00 mm) obtained by the rosemary EBr with that (16.62 mm) obtained by the ethanolic extract tested by Zhang *et al.*, [36].

The differences found can be attributed to several reasons such as inherent factors, methods of extraction ([8]; [23]; [37]), preparation of the extract, solvent used, the sensitivity of the bacteria [38], and finally the part of the plant used [31].

No inhibition zone was observed around disks impregnated of the different extracts. In the light of these results, we can conclude that the extracts from the two plants do not contain antifungal agent.

## CONCLUSION

Through this work, we want to show that the plants constitute a very interesting reservoir for research in the future. An extension of this work in the future is desirable to study the components present in the rosemary EAcOEt and to assess their antibacterial activities.

A search of the antibacterial agent responsible for the inhibition of the *Enterobacter* sp (ESBL+HLC) is also necessary and seems of great importance.

## REFERENCES

- [1]- SAGDIC O, KUSCU A, ÖZCAN M and ÖZCELİK S. Effects of Turkish spice extracts at various concentrations on the growth of *Escherichia coli* O157:H7. Food Microbiology. Vol 19. (2002). pp.473-480.
- [2]- BAHORUN T, Substances naturelles actives : la flore mauricienne, une source d'approvisionnement potentielle. Food and agricultural resarch council, Réduit, Mauritius. (1997). pp.83-94.
- [3]- ATIK BEKKARA F, BOUSMAHA L, TALEB BENDIAB S.A, BOTI J.B and CASANOVA J. Composition chimique de l'huile essentielle de *Rosmarinus officinalis* L poussant à l'état spontané et cultivé de la région de Tlemcen. Biologie & Santé. Vol 7. (2007). pp. 6-11.
- [4]- GONZALEZ-TRUJANO M.E, PENA E.I, MARTINEZ A.L, MORENO J, GUEVARA-FEFER P, DECIGA-CAMPOS M and LOPEZ-MUNOZ F.J. Evaluation of the antinociceptive effect of *Rosmarinus officinalis* L. using three different experimental models in rodents. J Ethnopharmacol. Vol 111.(2007) .pp. 476-482.
- [5]- WONG C.C, LI H.B, CHENG K.W and CHEN F. A systematic survey of antioxidant activity of 30 Chinese medicinal plants using the ferric reducing antioxidant power assay, Food Chem. Vol 97. (2006). pp.705-711.
- [6]- DJERIDANE A, YOUS M, NADJEMI B, BOUTASSOUNA D, STOCKER P and VIDAL N. Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. Food Chem. Vol 97. (2006). pp.654-660.
- [7]- AMAROWICZ R, TROSZYNsKA A and SHAHIDI F. Antioxidant activity of almond

seed extract and its fractions. *J food lipids.* Vol 12. (2005). pp.344-358.

[8]- CELIKTAS O.Y, HAMES KOCABAS E.E, BEDIR E, VARDAR SUKAN F, OZEK T and BASER K.H.C. Antimicrobial activities of methanol extracts and essential oils of *Rosmarinus officinalis*, depending on location and seasonal variations. *Food Chem.* Vol 100. (2007). pp.553-559.

[9]- NGAMENI B, KUETE V, SIMO I.K, MBAEVNG A.T, AWOUSSONG P.K., PATNAM R, ROY R and NGADJUI B.T. Antibacterial and antifungal activities of the crude extract and compounds from *Dorstenia turbinata* (Moraceae). *South African J Botany.* Vol 75. (2009). pp.256-261.

[10]- GORTZI O, LALAS S, CHINOU I and TSAKNIS J. Evaluation of the Antimicrobial and Antioxidant Activities of *Origanum dictamnus* Extracts before and after Encapsulation in Liposomes. *Molecules.* Vol 12. (2007). pp.932-945.

[11]- FECKA I and TUREK S. Determination of polyphenolic compounds in commercial herbal drugs and spices from Lamiaceae: thyme, wild thyme and sweet marjoram by chromatographic techniques. *Food Chem.* 108 (2008). pp.1039-1053.

[12]- ERKAN N, AYRANCI G and AYRANCI E. Antioxidant activities of rosemary (*Rosmarinus Officinalis* L.) extract, blackseed (*Nigella sativa* L.) essential oil, carnosic acid, rosmarinic acid and sesamol. *Food Chem.* Vol 110. (2008). pp.76-82.

[13]- HO S.C, TSAI T.H, TSAI P.J and LIN C.C. Protective capacities of certain spices against peroxynitrite-mediated biomolecular damage. *Food and Chemical Toxicology.* Vol 46 (2008). pp.920-928.

[14]- TSAI P, TSAI T and HO S. *In vitro* inhibitory effects of rosemary extracts on growth and glucosyltransferase activity of *Streptococcus sobrinus*. *Food Chem.*(2007) (in press).

[15]- TAWAHA K, ALALI F.Q, GHARAIBEH M, MOHAMMAD M and EL-ELIMAT T. Antioxidant activity and total phenolic content of selected Jordanian plant species. *Food Chem.* (2007) (in press).

[16]- GOMEZ-CARAVACA A.M, GOMEZ-ROMERO M, ARRAEZ-ROMAN D, SEGURA-CARRETERO A and FERNANDEZ-GUTIERREZ A. Advances in the analysis of phenolic compounds in products derived from bees. *J Pharmaceutical and Biomedical Analysis.* Vol 41. (2006). pp.1220-1234.

[17]- FALLEH H, KSOURI R, CHAIEB K, KARRY-BOURAQUI N, TRABELSI N, BOULAABA M and ABDELLY C. Phenolic composition of *Cynara cardunculus* L. organs, and their biological activities. *C. R. Biologies.* Vol 331. (2008). pp.372-379.

[18]- MAISUTHISAKUL P, PASUK S and RITTHIRUANGDEJ P. Relationship between antioxidant properties and chemical composition of some Thai plants. *J Food Composition and Analysis.* Vol 2.(2008). pp.229-240.

[19]- MERKEN H.M and BEECHER G.R. Liquid chromatographic method for the separation and quantification of prominent flavonoid aglycones. *J Chromatography A.* Vol 897. (2000). pp.177-184.

[20]- JUSTESEN U and KNUTHSEN P. Composition of flavonoids in fresh herbs and calculation of flavonoid intake by use of herbs in traditional Danish dishes. *Food Chemistry.* Vol 73. (2001). pp.245-250.

[21]- WOJDYLO A, OSZMIANSKI J and CZEMERYS R. Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chem.* Vol 105. (2007). pp.940-949.

[22]- HAYOUNI E.A, ABEDRABBA M, BOUIX M and HAMDI M. The effects of solvents and extraction method on the phenolic contents and biological activities *in vitro* of Tunisian *Quercus coccifera* L. and *Juniperus phoenicea* L. fruit extracts. *Food Chem.* (2007) (in press).

[23]- TURKMEN N, VELIOGLU Y. S, SARI F and POLAT G. Effect of Extraction Conditions on Measured Total Polyphenol Contents and Antioxidant and Antibacterial Activities of Black Tea. Molecules. Vol 12. (2007). pp.484-496.

[24]- SHAN B, CAI Y.Z, BROOKS J.D and CORKE H. The *in vitro* antibacterial activity of dietary spice and medicinal herb extracts. International J Food Microbiology. Vol 117. (2007). pp.112- 119.

[25]- KONÉ W. M, KAMANZI ATINDEHOU K, TERREAUX C, HOSTETTMANN K, TRAORÉ D and DOSSO M. Traditional medicine in North Côte-d'Ivoire: screening of 50 medicinal plants for antibacterial activity. J Ethnopharmacol. Vol 93. (2004). pp.43-49.

[26]- GEORGANTELIS D, AMBROSIADIS I, KATIKOU P, BLEKAS G and GEORGAKIS S.A. Effect of rosemary extract, chitosan and  $\alpha$ -tocopherol on microbiological parameters and lipid oxidation of fresh pork sausages stored at 4 °C. Meat Science. Vol 76. (2007). pp.172-181.

[27]- BAILENTINE C.W, CRANDALL P.G, O'BRYAN C.A, DUONG D.Q and POHLMAN F.W. The pre- and post-grinding application of rosemary and its effects on lipid oxidation and color during storage of ground beef. Meat Science. Vol 73. (2006). pp. 413-421.

[28]- GUESMI A et BOUDABOUS A. Activité antimicrobienne de cinq huiles essentielles associées dans les produits de thalassothérapie. Revue des Régions Arides. Numéro spécial (2006). pp.224-230.

[29]- DORDEVIC S, PETROVIC S, DOBRIC S, MILENKOVIC M, VUCICEVIC D, ZIZIC S and KUKIC J. Antimicrobial, anti-inflammatory, anti-ulcer and antioxidant activities of *Carlina acanthifolia* root essential oil. J Ethnopharmacol. Vol 109. (2007). pp.458 -463 .

[30]- RASOOLI I, FAKOOR M.H, YADEGARINIA D, GACHKAR L, ALLAMEH A. REZAEI M.B. Antimycotoxicogenic characteristics of *Rosmarinus officinalis* and *Trachyspermum copticum* L. essential oils. International J of Food Microbiology. Vol 122. (2008). pp.135-139.

[31]- NATARAJAN D, John Britto S, Srinivasan K, Nagamurugan N, Mohanasundari C and Perumal G. Anti-bacterial activity of Euphorbia fusiformis-A rare medicinal herb. J Ethnopharmacol. Vol 102. (2005). pp.123-126.

[32]- FAZELI M. R, AMIN G, AHMADIAN-ATTARI M. M, ASHTIANI H, JAMALIFAR H. and SAMADI N. Antimicrobial activities of Iranian sumac and avishan-e shirazi (*Zataria multiflora*) against some food-borne bacteria. Food Control. Vol 18. (2007). pp.646-649.

[33]- ASKUN T, TUMEN G, SATIL F and ATES M. *In vitro* activity of methanol extracts of plants used as spices against *Mycobacterium tuberculosis* and other bacteria. Food Chem. Vol 116. (2009). pp.289-294.

[34]- KIL H.Y, SEONG E.S, GHIMIRE B.K, CHUNG I.M, KWON S.S, GOH E.J, HEO K, KIM M.J, LIM J.D, LEE D and YU C.Y. Antioxidant and antimicrobial activities of crude sorghum extract. Food Chem. Vol 115. (2009). pp.1234-1239.

[35]- MOHSEN S. M and AMMAR A.S.M. Total phenolic contents and antioxidant activity of corn tassel extracts. Food Chem. Vol 112. (2009). pp. 595-598.

[36]- ZHANG H, KONG B, XIONG Y.L and SUN X. Antimicrobial activities of spice extracts against pathogenic and spoilage bacteria in modified atmosphere packaged fresh pork and vacuum packaged ham slices stored at 4 °C. Meat Science. Vol 81. (2009). pp. 686-692.

[37]- SAGDIC O and OZCAN M. Antibacterial activity of Turkish spice hydrosols. *Food Control*. Vol 14. (2003). pp.141-143.

[38]- LOZIENE K, VENSKUTONIS P. R, SIPAILIENÉ A and LABOKAS J. Radical scavenging and antibacterial properties of the extracts from different *Thymus pulegioides* L. chemotypes. *Food Chem*. Vol 103. (2007). pp.546-559.